



Steroid Withdrawal Effects Following Long-term Topical Corticosteroid Use.

Sheary B¹.

+ Author information

Abstract

BACKGROUND: Concerns about topical steroid withdrawal (TSW) are leading some patients to cease long-term topical corticosteroid (TCS) therapy. Diagnostic criteria for this condition do not exist.

OBJECTIVE: The aim of this study was to examine the demographics and outcomes in adult patients who believe they are experiencing TSW following discontinuation of chronic TCS overuse.

METHODS: This was a retrospective cohort study of patients in an Australian general practice presenting with this clinical scenario between January 2015 and February 2018.

RESULTS: Women represented 56% of the 55 patients seen, and ages ranged from 20 to 66 years (mean, 32.9 years; median, 30.0 years). Seventy-six percent had an original diagnosis of atopic dermatitis. Sixty percent had used potent TCSs on their face, and 42% had a history of oral corticosteroid use for skin symptoms. Burning pain was reported in 65%, all had widespread areas of red skin, and so-called "elephant wrinkles," "red sleeve," and the headlight sign were seen in 56%, 40%, and 29%, respectively.

CONCLUSIONS: Patients with a history of long-term TCS overuse may experience symptoms and signs described in TSW on stopping TCSs. Diagnostic criteria, reflecting the histories and examination findings of the patients studied, are suggested in this article with the aim to advance discussion and research into TSW. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

